

Name: \_\_\_\_\_ ID \_\_\_\_\_

**Posterior Distributions and Statistical Inference.**

1. A study comparing coffee drinkers with non-drinkers found that the relative risk (RR) of pancreatic cancer had  $p$ -value = .15. Indicate with an “X” which of the following is consistent with that  $p$ -value.

X	95% CI for RR	Significant?
<input type="checkbox"/>	85 to 3.24	Significant
<input type="checkbox"/>	0.85 to 3.24	Not significant
<input type="checkbox"/>	1.15 to 3.01	Significant
<input type="checkbox"/>	1.15 to 3.01	Not significant
<input type="checkbox"/>	-.53 to .67	Not significant

2. Two hundred forty-four male alcoholics suffering from secondary hypertension participated in a study to determine the efficacy of a new antihypertensive agent. The men were assigned at random to either the control group or the treatment group. Men in the control group received a placebo. Statistics for arterial pressure at 30 days after treatment are shown in this table.

<i>Hypertension Study</i>	Placebo	Treatment
n	100	144
mean ( $\bar{x}$ )	140.3	102
standard deviation (s)	15.0	7.2

Describe the approximate posterior distribution of the difference ( $\Delta$ ) between the means of the two populations and compute a 95% confidence interval for the difference.

The posterior distribution is: \_\_\_\_\_

\_\_\_\_\_

95% Confidence interval: \_\_\_\_\_ to \_\_\_\_\_

3. In a study of paranoid schizophrenics, the age of onset of each person’s illness was determined. Statistics for age at onset are shown in this Table.

<i>Schizophrenia Study</i>	Males	Females
n	9	12
mean ( $\bar{x}$ )	26.56	29.58
standard deviation (s)	29.78	38.45

Describe the approximate posterior distribution of the difference ( $\Delta$ ) between the means of the two populations and compute a 95% confidence interval for the difference.

The posterior distribution is: \_\_\_\_\_

\_\_\_\_\_

95% Confidence interval: \_\_\_\_\_ to \_\_\_\_\_

Is the difference statistically significant?: No  Yes  What tells you this?: \_\_\_\_\_

4. A study by Mann et al. investigated whether oral contraceptive use increased the rate of heart attacks. The data are in this table,

<b>Contraceptive Study</b>	Used Oral Contraceptives	Never Used Oral Contraceptives
n	57	167
% having heart attacks	23%	35%

Compute the proportions having heart attacks. Describe the posterior distribution of  $\Delta$ , the difference between the population proportions. Obtain a 95% confidence interval for the difference between the population proportions. Is the difference statistically significant? Explain why or why not.

a) Posterior Distribution: \_\_\_\_\_

b) 95% Confidence Interval: \_\_\_\_\_

c) Statistically Significant?: Yes  No  Why?: \_\_\_\_\_

5. One possible side effect of antihistamines is drowsiness. In a clinical trial of the drug Suspirizine vs. a placebo. The relative risk of drowsiness in the drug group relative to the placebo group was  $RR = (8\%/6\%) = 1.33$  and the 95% CI was 1.09 to 1.69. Presumably if a drug produces a “very important” increase in the risk of a side effect the FDA will require a warning label; however we don't know how big the RR has to be to be considered “very important,” so let's work two scenarios.

a) The FDA thinks  $RR > 1.5$  is very important. Compute the probability. What do you conclude from the study?

b) The FDA thinks  $RR > 2.0$  is very important. Compute the probability. What do you conclude from the study?

### Statistical Models

6. The patients in the data set in the table below are wearing one of two different cochlear implant devices. Suppose I want a regression model with different intercepts but the same slope for the two devices. In the following spreadsheet fill in columns X1, X2, and X3 with the appropriate design variables so that  $\beta_1$  is the intercept for device A,  $\beta_2$  is the intercept for device B, and  $\beta_3$  is the common slope for the two devices.

Data			Design Variables		
Device	Month 1 (x)	Asymptotic (y)	X1	X2	X3
A	20	31.00			
A	35	70.33			
A	26	48.75			
...	...	...	...	...	...
B	0	25.50			
B	22	44.00			
B	29	40.00			
...	...	...	...	...	...

7. A statistician computed a multiple linear regression model for the relationship between salary (y), race, gender, and years of experience. This table shows the regression coefficients, their standard errors, p-values, and confidence intervals.

Variable	$\hat{\beta}$	SE	p-value	95% Confidence Interval	
				Lower	Upper
Intercept	30,000	5,000	.0001	20,000	40,000
Male	400	180		40	760
White	700	250			
Years	500	60	.001	300	500

What is the average salary for **black males** with 5 years of experience? \_\_\_\_\_

What is the average salary for **white males** with 5 years of experience? \_\_\_\_\_

Find a 95% confidence interval for the difference \_\_\_\_\_ to \_\_\_\_\_

Is the difference significant? \_\_\_\_\_

What tells you this? \_\_\_\_\_

8. The table below shows the results of a (fictional) study of 10-year survival for men who survived a heart attack. The outcome variable is surviving 10 years after first heart attack (1=survived / 0=died), so the logistic regression model predicts the logarithm of the odds on survival. The explanatory variables are family history of early death (1 = at least one first degree male relative died before the age of 55 / 0 = none), body mass index, and age at the time of the first heart attack.

Variable	$\hat{\beta}$	SE	odds ratio	95% Confidence Interval for odds ratio	
				Lower	Upper
1 Intercept	8.1548	0.8864	na	na	na
2 Family History	-0.0513	0.0112			
3 Body Mass Index	-0.1020	0.1627	0.903	0.656	1.242
4 Age	-0.1069	0.0220	1.113	1.066	1.162

**Table 1.** Logistic regression coefficients for 10-year survival of males following first heart attack.

Compute the estimated probability of survival for a 58-year-old with a family history of early death and BMI = 30 .

log(odds): \_\_\_\_\_ odds: \_\_\_\_\_ prob: \_\_\_\_\_

Which of the explanatory variables is(are) not statistically significant? What tells you this?

Insignificant variable(s): \_\_\_\_\_

What tells this: \_\_\_\_\_

What is the odds ratio for patients with a family history of early death vs. patients without a family history of early death?

Odds ratio: \_\_\_\_\_

Compute a 95% confidence interval for  $\beta_2$  (family history of early death) and for  $\exp(\beta_2)$ .

Confidence interval for  $\beta_2$ : \_\_\_\_\_ to \_\_\_\_\_

Confidence interval for  $\exp(\beta_2)$ : \_\_\_\_\_ to \_\_\_\_\_

9. A study was conducted to determine if maternal smoking influenced the birth weight of infants. Birth weight (pounds) is also influenced by gestational age (number of days between conception and birth). Design variables were the intercept, whether the mother smoked during pregnancy (1=yes/0=no), and the gestational age of the infant (days). The regression results were as follows:

Variable	$\hat{\beta}$	Std Err	95% CI	
			Lower	Upper
Intercept	-1.68	130		
Gestational Age	.052	.022		
Mother Smoked	-.25	.09		

Compute the confidence intervals for the regression coefficients. Does maternal smoking have a significant effect on birth weight? How do you know this?

Significant (Y/N)? \_\_\_\_\_ Because: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

10. If the credible interval for a logistic regression coefficient ( $\beta$ ) rules out zero, what can you say about a) the credible interval for the odds ratio, and b) the  $p$ -value?

\_\_\_\_\_


\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

11. The diagram below is a decision problem in tree form. A patient has been exposed to a disease and may or may not be infected but is not yet manifesting symptoms. The physician can call for a diagnostic test or not. The test costs \$7 and is not perfect. Based on the test results (or not) the physician can decide to prescribe a \$35 prophylactic medication that prevents the disease if it is applied before symptoms appear. If the physician waits for symptoms to appear, it is necessary to prescribe a rescue medication that costs \$100. The prevalence of the disease is 0.2 and the test has sensitivity .98 and specificity .999. Solve the decision tree.




12. The next table shows the calculation of the Kaplan-Meier survival function for these survival times:

1 2 5 6 7 9 15 15 >21 21 >25 >30 31 39 40 >40

Interval	at Risk	Failed	Censored	Failure Rate	Survival Rate
0	n/a	n/a	n/a	n/a	1.0000
0 - 1	16	1	0	0.0625	0.9375
1 - 2	15	1	0	0.0667	0.8750
2 - 5	14	1	0	0.0714	0.8166
5 - 6	13	1	0	0.0769	<b>C</b>
6 - 7	12	1	0	0.0833	0.6875
7 - 9	11	1	0	0.0909	0.6250
9 - 15	10	2	0	0.2000	0.5000
15 - 21	8	1	1	<b>B</b>	0.4375
21 - 25	<b>A</b>	0	1	0.0000	0.4375
25 - 30	5	0	1	0.0000	0.4375
30 - 31	4	1	0	0.2500	0.3281
31 - 39	3	1	0	0.3333	0.2188
39 - 40	2	1	1	0.5000	0.1094

Fill in the missing numbers for cells A, B, and C.

13. The following graph shows Kaplan-Meier survival curves for three groups of patients (I, II, and III). Time is expressed in months.



How do the groups compare in terms of survival time, from longest to shortest?

- a) III>II>I      b) I>II>III      c) II>III>I      d) II>I>III      e) Impossible to determine.

What is the median survival time in Group II? (Draw it on the graph).

What is the 100-month survival rate for patients in group I? (Draw it on the graph.)

14. A sample of  $n=400$  drawn from a fairly normal population has sample mean  $\bar{x} = 16.5$  and sample standard deviation  $s = 8.8$ . Describe the approximate posterior distribution of the population mean  $\mu$  and obtain an approximate 95% confidence interval for  $\mu$ .

The posterior distribution is: \_\_\_\_\_

95% Confidence interval: \_\_\_\_\_ to \_\_\_\_\_

*(more questions on next page)*

15. An unknown quantity  $\Delta$  has a  $t(11)$  distribution with  $\mu = 20$  and  $\sigma = 5$ . Compute the 95% credible interval for  $\Delta$ . Fill in the blanks in the `ttailarea.xls` spreadsheet to compute  $P(\Delta > 24)$

<b>Instructions:</b>		
Enter degrees of freedom and z (or t) value.		
Read left- or right tail area.		
Tail areas for t(df)		
	v =df =	
	t=z =	
Left Tail	$P(t(df) \leq z) =$	
Right Tail	$P(t(df) > z) =$	

Enter Here

Read Here