

- problem: to find values of variable x that satisfy $f(x) = 0$ for given function f
- solution is called “zero of f ” or “root of f ”
- when is this an important problem in statistics?

The bisection method

- also called “binary-search method”
- conditions for use
 - f continuous, defined on interval $[a, b]$
 - $f(a)$ and $f(b)$ of opposite sign
- by Intermediate Value Theorem, there exists a p , $a < p < b$, such that $f(p) = 0$
- procedure works when $f(a)$ and $f(b)$ of opposite sign and more than one root in $[a, b]$
- for simplicity, we’ll assume unique root in interval
- method consists of
 - repeated halving of subintervals of $[a, b]$
 - at each step, locating half containing p
- requires following inputs
 - endpoints a, b
 - tolerance TOL
 - maximum number of iterations N_0

Fixed-point iteration

- solution to $g(x) = x$ is called *fixed point* of function g
- Theorem
 - conditions
 - * g continuous on $[a, b]$
 - * $g(x) \in [a, b] \forall x \in [a, b]$
 - conclusions
 - * g has a fixed point in $[a, b]$
 - if further
 - * $g'(x)$ exists on (a, b) and a positive constant $k < 1$ exists such that $|g'(x)| \leq k < 1, \forall x \in (a, b)$
 - then
 - * g has a unique fixed point p in $[a, b]$

Example 1

$$g(x) = \frac{x^2 - 1}{3} \text{ on } [-1, 1]$$

- absolute minimum of g is $g(0) = -\frac{1}{3}$
- absolute maximum of g is $g(\pm 1) = 0$
- $|g'(x)| = |\frac{2x}{3}| \leq \frac{2}{3} \forall x \in [-1, 1]$
- so g has unique fixed point p in interval
- in this case, can be determined exactly by quadratic formula

Example 2

$$g(x) = 3^{-x} \text{ on } [0, 1]$$

- $g(1) = \frac{1}{3} \leq g(x) \leq 1 = g(0) \forall 0 \leq x \leq 1$, so fixed point exists in interval
- theorem cannot be used to determine uniqueness of fixed point since $|g'(0)| = 1.0986 > 1$
- but fixed point must be unique since g is a decreasing function

5

Fixed-Point Theorem

- conditions
 - g continuous on $[a, b]$
 - $g(x) \in [a, b] \forall x \in [a, b]$
 - $g'(x)$ exists on (a, b) and

$$|g'(x)| \leq k < 1, \forall x \in (a, b)$$
- then if p_0 is any number in $[a, b]$ then the sequence defined by

$$p_n = g(p_{n-1}), \quad n \geq 1$$
 converges to the unique fixed point p in $[a, b]$

7

Fixed-point iteration

- choose initial approximation p_0
- set $p_n = g(p_{n-1})$ for each $n \geq 1$

Example

$x^3 + 4x^2 - 10 = 0$ has unique root in $[1, 2]$.

6

The Newton-Raphson Method

- one of most powerful and well-known numerical methods for solving root-finding problem $f(x) = 0$
- one derivation: Taylor series approximation
 - suppose f' and f'' are continuous on $[a, b]$
 - let $x_0 \in [a, b]$ be an approximation to p such that $f'(x_0) \neq 0$ and $|x_0 - p|$ is “small”
 - first order Taylor approximation for $f(x)$ expanded around x_0

$$f(x) = f(x_0) + (x - x_0)f'(x_0) + \frac{(x - x_0)^2}{2}f''(\xi(x))$$
 where $\xi(x)$ is between x and x_0 .
 - with $x = p$ this gives

$$0 = f(x_0) + (p - x_0)f'(x_0) + \frac{(p - x_0)^2}{2}f''(\xi(x))$$
 - since $|x_0 - p|$ is “small”, $(x_0 - p)^2$ should be negligible and

$$0 \simeq f(x_0) + (p - x_0)f'(x_0)$$
 - solving for p yields

$$p \simeq x_0 - \frac{f(x_0)}{f'(x_0)}$$

8

The Newton-Raphson Method

- start with initial approximation p_0
- let $p_n = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})}$

Convergence Theorem for Newton-Raphson Method

- conditions
 - f has continuous first and second derivatives on $[a, b]$
 - $p \in [a, b]$ is such that $f(p) = 0$ and $f'(p) \neq 0$
- conclusions
 - then there exists a $\delta > 0$ such that Newton's method generates a sequence $\{p_n\}_{n=1}^{\infty}$ converging to p for any initial approximation $p_0 \in [p - \delta, p + \delta]$.

The Secant Algorithm

- useful when computation of $f'(x)$ is far more computationally intensive than computation of $f(x)$
- uses forward (or backward)-difference formula to approximate $f'(p_{n-1})$

$$f'(p_{n-1}) \simeq \frac{f(p_{n-2}) - f(p_{n-1})}{p_{n-2} - p_{n-1}} = \frac{f(p_{n-1}) - f(p_{n-2})}{p_{n-1} - p_{n-2}}$$

- Secant algorithm generates sequence as

$$p_n = p_{n-1} - \frac{f(p_{n-1})(p_{n-1} - p_{n-2})}{f(p_{n-1}) - f(p_{n-2})}, \quad n \geq 1$$