

**22S:30/105**  
**Statistical Methods and Computing**

**Inference for Proportions, continued**

Lecture 19  
 Apr. 7, 2008

Kate Cowles  
 374 SH, 335-0727  
 kcowles@stat.uiowa.edu

**Hypotheses**

The null hypothesis says that the population proportion  $p$  in those diagnosed before age 40 is the same as the known proportion in those diagnosed at a later age.

$$H_0 : p = 0.082$$

The alternative hypothesis is two-sided because we do not know in advance in which direction a difference might go. (Younger people in general are more likely to survive for 5 years than older people, but perhaps a more severe form of lung cancer occurs in younger people.)

$$H_a : p \neq 0.082$$

**Significance level**

We choose to do our test at the  $\alpha = .05$  significance level.

**Single-sample hypothesis testing about a proportion**

Example:

- We know from large databases of medical records that, among patients diagnosed with lung cancer when they are 40 years of age or older, the proportion that survive for 5 years after diagnosis is 0.082.
- We are interested in determining whether the proportion of 5-year survivors is the same in the population of patients diagnosed with lung cancer before age 40.
- The parameter of interest is the population proportion  $p$  in the population diagnosed with lung cancer before age 40.
- We will get data on a sample of persons under 40 who have been diagnosed with lung cancer.

**Data**

From a 1991 article in the journal *Cancer*, we obtain data on a sample of 52 person diagnosed with lung cancer at age 40 or younger. Only 6 of them survived for 5 years after diagnosis.

The sample proportion was

$$\hat{p} = \frac{6}{52} = 0.115$$

**The test statistic**

The  $z$  test statistic is:

$$\begin{aligned} z &= \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}} \\ &= \frac{0.115 - 0.082}{\sqrt{\frac{0.082(1-0.082)}{52}}} \\ &= 0.87 \end{aligned}$$

## The p-value

Because the test is two-sided, the p-value is the area under the standard normal curve more than 0.87 away from 0 *in either direction*. Table A tells us that the area to the left of -0.87 is 0.192. The p-value is twice this area:

$$p = 2(0.192) = 0.384$$

**The 95% confidence interval for the proportion  $p$  of patients diagnosed with lung cancer before age 40 who will survive 5 years is:**

$$\begin{aligned} \hat{p} \pm z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} &= 0.115 \pm 1.96 \sqrt{\frac{(0.115)(1-0.115)}{52}} \\ &= 0.115 \pm 0.087 \\ &= (0.028, 0.202) \end{aligned}$$

## Conclusion

Can we reject the null hypothesis that  $p = 0.082$ ?

A proportion of survivors as far from 0.082 as what we found would happen 38% of the time if a sample of 52 patients were drawn from a population in which the true proportion of survivors was 0.082. Our result does not show that the proportion of 5-year lung cancer survivors is different in the population of patients diagnosed before age 40 from in the population diagnosed at age 40 or later.

**Choosing the sample size for a desired margin of error**

- Recall that the **margin of error** is the quantity that we add to and subtract from a point estimate in order to compute the right and left endpoints of a confidence interval.
- For a proportion, the confidence interval is

$$\hat{p} \pm z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

- so the margin of error is

$$z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

- Since we don't know in advance what  $\hat{p}$  is going to be, we have to guess it. Call our guess  $p^*$ . Some ways to make an "educated guess":
  - Use a pilot study or past experience with similar studies.
  - Use  $p^* = 0.5$ . This is conservative, since it will give the largest possible margin of error.
- Then if  $m$  is the desired margin of error, the required sample size  $n$  is:

$$n = \left(\frac{z^*}{m}\right)^2 p^*(1 - p^*)$$

- How would the sample size change if you had no previous information about what proportion to expect?

### Example:

- PTC is a substance that has a strong bitter taste for some people and is tasteless for others. The ability to taste PTC is inherited. About 75% of Italians can taste PTC, for example.
- You want to estimate the proportion of Americans with at least one Italian grandparent who can taste PTC.
- Starting with the 75% estimate for Italians, how large a sample must you test in order to estimate the proportion of PTC tasters within  $\pm 0.04$  with 95% confidence?

### Sample size calculation for a hypothesis test regarding a single population proportion

- Consider a one-sided test:

$$H_0 : p = p_0$$

$$H_a : p < p_0$$

- To compute sample size, we need to specify:
  - the significance level  $\alpha$
  - a specific alternative hypothesis  $p = p_1$
  - the power  $1 - \beta$
- Then the sample size  $n$  is

$$n = \left[ \frac{z_{1-\alpha}\sqrt{p_0(1-p_0)} + z_{1-\beta}\sqrt{p_1(1-p_1)}}{(p_1 - p_0)} \right]^2$$

**Example:**

- Suppose in the PTC example that instead of just estimating  $p$  in Americans with at least one Italian grandparent, we wished to test the hypotheses:

$$H_0 : p = .75$$

$$H_a : p < .75$$

- We choose
  - $\alpha = .05$
  - We would not consider the difference to be scientifically meaningful unless the true  $p$  were .60 or less, so we set  $p_1 = .6$ .
  - We want 90% power if the true  $p$  is .6.

- According to Table A

$$- z_{1-\alpha} = 1.645$$

$$- z_{1-\beta} = 1.28$$

- So our sample size is

$$\begin{aligned} n &= \left[ \frac{1.645\sqrt{.75(.25)} + 1.28\sqrt{.6(.4)}}{(.6 - .75)} \right]^2 \\ &= 8.929^2 \\ &= 79.73 \text{ or } 80 \end{aligned}$$

For a two-sided test, use  $z_{1-\frac{\alpha}{2}}$  instead of  $z_{1-\alpha}$  in the formula:

$$n = \left[ \frac{z_{1-\frac{\alpha}{2}}\sqrt{p_0(1-p_0)} + z_{1-\beta}\sqrt{p_1(1-p_1)}}{(p_1 - p_0)} \right]^2$$

In our example, this would be:

$$\begin{aligned} n &= \left[ \frac{1.96\sqrt{.75(.25)} + 1.28\sqrt{.6(.4)}}{(.6 - .75)} \right]^2 \\ &= 9.838^2 \\ &= 96.8 \text{ or } 97 \end{aligned}$$